



13 Rules

On Preventing and Controlling African Swine Fever within Farms

PREVENTION & CONTROL



1 Raise awareness. Enhance training on the prevention and containment of African swine fever and perform tasks strictly in compliance with anti-epidemic procedures.



2 Isolate the farm. Set up a "No visit during outbreak" sign 100m away from the farm. Strictly restrict the number of persons entering the farm, allowing entry only when necessary.

SURROUNDINGS / VISITORS



3 Build a **pig transfer center** or a loading platform 3 km away from the farm. Prevent direct contact of farms with external pig transportation vehicles to reduce the risk of transfer of infection from truck to the farm.

CLEANLINESS & DISINFECTION



4 Biosecure disposal system. In accordance with national regulations, set up channels specific for deceased pigs and fecal matter to ensure disposal that will not spread the disease. Such channels must not be linked to other channels.



5 Institute **internal and external cleaning and disinfection stations.** The internal cleaning and disinfection station is exclusively used for feed and pig transportation vehicles and staff vehicles inside the farm. The external cleaning and disinfection station is set up 3 km away from the farm and used for external feed transportation vehicles, external staff vehicles and pig collection vehicles.



6 For all vehicles, institute a **5-step cleaning and disinfection procedure** as follows: (1) Debris removal through water spraying, (2) foaming (use of surfactants), (3) washing, (4) disinfection and (5) drying.

FEED SAFETY & FOOD SAFETY



7 Implement feed hygiene. Choose feed from reliable brands to minimize cross-infection risk and ensure traceability. During transport, feeds should be in an enclosed van or protected by a cover. In going to the farm, disinfected feed trucks should use a separate road different from the road where pig trucks pass. External feed transportation vehicles are not allowed to enter the farm. Unloading and transfer of feed to an internal farm vehicle is encouraged.



8 Disinfect external purchases of food and other consumables. All items purchased externally must be disinfected through ozone fumigation before being brought into the farm.



9 No to swill feeding. Swill feed may contain virus-contaminated meat and water.



10 No external pork and pork products. Prohibit the purchase or entry of fresh, cooked or preserved pork products into the farm.

ANIMAL HANDLING

Logistics
Dead animal processing
Animals and Insects



11 Closed farm policy. Do not purchase any breeding pigs or semen from external sources. If necessary, purchase breeding stocks or semen only from ASF free certified farms.



12 Animal control. Do not keep cats, dogs, livestock or other pets in the farm.



13 Bird and rodent control. Set up bird netting, fly netting and mosquito netting, and perform rodent extermination within the grounds on a regular basis.

